

File attachment #101
Mixing AutoMetrics with MARS: A study of
psychological sequelae of Chernobyl
(Hypotheses 4, 6)

DRU: Modeling Nuclear Disaster Risk: The Effects of Perceived Risk and Radiation
Exposure on Post-Chernobyl Psychosocial and Health Behavior Outcomes in
Ukrainian Residents
NSF Grant 082 6983

30 September 2012

Contents

1	Acknowledgements	2
2	Introduction	2
3	Nomenclature	4
4	Data collection and sampling	5
5	Research instruments and measures	6
5.1	Outcome measures	6
5.1.1	Physical health	6
5.2	Outcome measures	7
5.2.1	Mental health	7
5.2.2	Predictors	8
6	Objectives	9
7	Historical highlights in the development of MARS	12
8	MARS algorithm	14
8.1	How MARS works	14
8.2	basis function generation	17
9	Dose reconstruction of exposure to ¹³⁷CS	22
10	Application of MARS and AutoMetrics to variable selection and model building	26
10.1	BSI positive symptoms models	26
10.1.1	Female positive symptoms models	26
10.1.2	Female positive symptoms model using AutoMetrics alone	26
10.1.3	Female model using AutoMetrics and MARS	27
10.2	Male Models for BSI Positive symptoms	28
10.2.1	Male model with AutoMetrics Alone	29
10.2.2	Male model for Positive symptoms using AutoMetrics + MARS	30
10.3	PTSD models	31
10.4	Male PTSD models	32
10.4.1	Male PTSD model using AutoMetrics alone	35
10.4.2	Male PTSD model using AutoMetrics + MARS	36
10.5	Male PTSD model with AutoMetrics + MARS- cont'd	37
10.6	Female PTSD models using AutoMetrics alone	42
10.6.1	Female PTSD model using AutoMetrics and MARS	48

11 Conclusion	52
11.1 Exploration of new dimensions	52
11.2 Are MARS models well-behaved?	53
11.3 Can MARS capture nonlinear relationships ?	53
11.4 Content validity Tables	54
11.5 Advantages of MARS	57
11.6 Disadvantages of MARS	58
11.7 Caveats	59

1 Acknowledgements

The project was funded by the National Science Foundation, Division of Decision, Risk and Uncertainty (082-6983); and conducted in cooperation with the Ministry of Health of Ukraine. HSD grant 08262983 and we are very grateful for their support. We would also like to thank the Ministry of Health in the Ukraine for their cooperation in this study. We are grateful to the developers of Stata in College Station, Texas, Salford Systems, Inc. in San Diego, California, and all of the AutoMetrics developers in Oxford, London for the software they have developed and which have used for variable selection and model building on this project.

2 Introduction

The current study conducted a random population sampling of Ukrainian residents in the Kiev and Zhytomir oblasts of that country, with the aim of developing long-term models of human nuclear disaster risk. Living in relatively close proximity to the Chornobyl Nuclear station in Ukraine, residents were exposed to the largest industrial radiological accident to date in 1986. A survey methodology was used to assess the complex bio-psycho-social pathways that contribute to long-term population outcomes after a significant radiological event. Data collection was conducted from 2008-2011.

In the effort to understand the long-term burden of nuclear accident exposure on a general population, investigators primary foci of interest included: the populations reconstructed cumulative dose exposure to ¹³⁷ Caesium (radiation source term for the Chornobyl event), cognitive perception of risk to health and environment, mental health status (standardized instruments), medical diagnoses (ICD-9), psychosocial functioning, health behavior, reproductive patterns, nutritional practices, Chornobyl accident information sources, and social communication networks. These domains were assessed in the population for their current status, and retrospectively for three earlier time periods from 1986 to the time of the survey interview. The current presentation describes preliminary statistical exploration of covariates influencing: 1) health-related behaviors as measured by the Nottingham Health Profile, 2) general mental health dysfunction as measured by the Brief Symptom Inventory , and 3) Chornobyl-related

post traumatic stress syndrome as measured by the Revised Civilian PTSD Scale.

Research of protracted, low dose radiation exposures from nuclear plant accidents, show relatively low impacts on population health risk [22]. Conversely, population perceptions of risk from radiological events remain active up to several years post-event [1, 3, 5, 14]. Toxic accidents (radiological, chemical, biologic agents) appear to drive a unique spectrum of psychosocial and behavioral responses in affected populations. These responses are distinct from event-related physical injury. They include depression, anxiety, traumatic response, increased medical services utilization, phobic nutritional behavior and changes in reproductive patterns [2, 3, 5, 6, 7].

3 Nomenclature

Multivariate adaptive regression splines (MARS)

- MARS is a method of using regression splines called basis functions to model local slope and intercept changes in relationships.
- It is similar to a piecewise regression model in that it combines basis functions to approximate the global nonlinear form of the relationship being modeled.
- MARS endeavors to improve the fit of the model and can identify interactions among the variables selected. To avoid overfitting, it uses a backward stepwise algorithm to trim the model to what it deems the essentials.

4 Data collection and sampling

- A sampling of 702 participants
- Randomized selection of phone numbers
- household sample of the Kiev and Zhitomir oblasts (states) of Ukraine
- Informed and consenting respondents were interviewed
- Responses were entered in interviewers hand-held computers,
- uploaded for storage to a website constructed for the study (Vovici Corporation, USA).
- Participants ranged in age from 28-84.

5 Research instruments and measures

5.1 Outcome measures

5.1.1 Physical health

- ICD-9 Medical Diagnosis. Diagnostic information from the Ukraine Ministry of Health database of annual standardized dispensary exams.
- Randomized selection of phone numbers
- Nottingham Health Profile. Standardized scale of self-reported health and its impact on multi-domain behavioral functioning [11]
- Reliability/validity on Russian language form tested in pilot study [12].

5.2 Outcome measures

5.2.1 Mental health

- Brief Symptom Inventory [13] Standardized scale measuring patterns of psychological distress: depression, anxiety, somatization, obsessive-compulsiveness, hostility, paranoia, psychoticism , global distress, positive symptom score. Russian form pilot tested [12].
- Revised Civilian PTSD Scale. Russian version anchored in Chornobyl event and restandardized [14]. Assesses post traumatic stress and distress clusters related to Chornobyl. Reliabilities computed in pilot study of this sample [12].
- of Psychosocial/Health Behavior: factors associated with toxic accident behavioral outcomes and are integrated within the research questionnaire, including medical services utilization, phobic behavior, substance use , eating habits, abortions and contraceptive use.

5.2.2 Predictors

- demographics,
- perception of radiation risk and nuclear attitudes, Chernobyl cognitions [1, 8],
- Chernobyl information sources, accident characteristics (distance, relocation, etc.),
- general hazards perception,
- negative life events and buffers,
- coping style [7].
- cumulative external radiation exposure to $^{137}\text{Caesium}$ was estimated for each participant (see below)

6 Objectives

Primary research objectives

- To explore the psychological sequelae of the Chernobyl disaster with respect to its effect on the part Ukrainian residents in the area.
- To analyze dose - mental health effect on survivors and residents of the general area.
- To analyze perceived risk of exposure effect on survivors and residents of the general vicinity of such a nuclear event.

Primary methodological objectives

- Although we use AutoMetrics for variable selection, we want to ascertain whether MARS can add value to this process.
- To answer two questions concerning the value added to our AutoMetrics analysis by MARS.

Does MARS help us explore dimensions of our data which we might have ignored?

Does MARS aid in providing a well-behaved model for our investigation of the subject of choice?

Could MARS help us capture the nonlinear relationships between variables?

Secondary objectives

- To use general-to-specific (GETS) approach to minimize omitted variable bias [10].
- To assure ourselves of not having overlooked important relevant variables.
- To be able to search, identify, and model non-linear functional relationships as painlessly as possible (while avoiding the curses of multidimensionality and multi- collinearity).
- We had a large number of variables– some of our datasets have approximately 3000 variables but we only have 339 males and 363 females in it.
- We could have more variables than observations in some cases and this is a problem *AutoMetrics* was designed to solve.
- To optimize construct validity– capturing all of the relevant and important dimensions
- To discover what disadvantages this usage entailed, if any.

7 Historical highlights in the development of MARS

- Automatic Interaction Detection: AID was developed by Morgan and Sonquist in 1963 in JASA to identify interaction terms by testing relative reduction in SSE. But there was a bias in the variable selection as the splits contained less observations.
- CHi-square Automatic Interaction Detection (CHAID) proposed by Gordon Kass in his doctoral dissertation in 1980 with bonferroni corrections for multiple tests.
- CART used regression and classification (Classification and regression trees) were later developed (Breiman, Friedman, Olshen and Stone, 1984) to combine categorical and continuous variables in the same tree output.
- Hastie, T. and Tibshirani, R. (1990) develop the generalized additive model which combines parametric and nonparametric estimation.
- Jerry Friedman developed MARS in 1993 as a method of automatic regression models— using basis functions and interactions between them.

Our situation

- Large datasets with more than 2900 variables in them.
- Sample size 363 women and 339 men
- Most analysis wrt socio-medical research is performed with gender-specific models.
- We want to developing optimal models to explain psychological phenomena experienced and reported by residents in the Oblasts near Chornobyl
- We have variables that exhibit a latency period before exhibiting a relationship
- We have variables that exhibit a delay manifesting a relationship.
- We depend on the multi-path misspecification testing of AutoMetrics to show guide our analysis.

8 MARS algorithm

8.1 How MARS works

MARS automates variable selection. It transforms variables into truncated regression splines to accommodate nonlinear functional forms. It checks for potential interactions with predictors. It drops irrelevant variables to minimize overfitting [23, 2].

MARS models nonlinear components with a variation of piecewise spline regression models. It uses a recursive partitioning algorithm for the regression problem, which emerged in a program for classification and regression trees, dubbed CART (Brieman, L., Friedman, J., Olshen, R. and Stone, C., 1984).

Originally MARS recursively partitioned the domain into disjoint subregions, within each of which it split the domain into two left and right offspring regions. The regression components would reside either within the left or the right daughter domain. Basis functions have the general form of

$$f(x) = \beta_0 + \sum_{j=1}^S h(m)B(m) \quad (1)$$

where $B(x) = I(x \in R_j)$ with I being an indicator function with a value of 1 if x is a condition is true or 0, otherwise. The indicator function then becomes part of a product of a series of step functions, in turn indicating membership of the subregion, formulated as

$$\begin{aligned} H[\eta] &= 1, \text{ if } \eta \geq 0, \\ &= 0, \text{ otherwise.} \end{aligned} \tag{2}$$

such that they could be incorporated as follows [19]

$$B_5(x) = H[x_1 - 0] * H[1 - x_2] * H[x_2 - 0] * H[1 - x_1], \tag{3}$$

which would delineate region 5 as a unit square within the larger region R . But as Stevens and Lewis point out these functions were step functions in disjointed areas. The disjoint requirement precluded MARS from finding linear additive models. Friedman replaced the univariate step functions with truncated regression splines.

A regression spline:

- If $x = 2\ 4\ 6\ 8\ 10\ 12\ 14\ 16\ 18\ 20$
- Spline formulation $(x-5)_+ = 0, 0, 0, 0, 0, 12, 14, 16, 18, 20$
- This spline may appear to be a shortened hockey stick with a value of zero until the 6th value is reached, upon the spline contains the values of x .

He formulated these in pairs, consisting of a primary basis function and a mirror image of it as

$$C = \{(X_j - k_t)_+, (k_t - X_j)_+\} \quad (4)$$

where k_t = represented the knots (joints) connecting the regression splines. The plus indicates that only the positive portion is used, so that the otherwise negative values are coded as zeros. This usually gives the basis function of the left-hand component with the parenthesis of Equation 5 the appearance of a hockey-stick. In effect, these functions entail recentering the variable around a knot. The effect of the transformation can take a variable that has a hockey - stick- like shape and linearize it, as shown in Figure 1.

8.2 basis function generation

- a stepwise algorithm. MARS begins with a constant and then searches each variable,
- and for each variable, tests all possible knots.
- The variable-knot combination that reduces the sum of squared errors the most or increases the sum of squared errors the least is selected.

Basis function transformation

- basis functions can transform hockey-stick like variables into more linear variables.
- For each primary transformation a mirror image is generated as well. For example if variable x is to be transformed, the basis function transformation might be $\max(0, 30 - x)$ and its mirror image would be $\max(0, x - 30)$.

What kinds of variables are amenable to such transformations?

- delayed responses or
- variables containing threshold conditions for a response to appear
- One example is the relationship between PTSD and anxiety shown in the left panel Figure 1, where a nonlinear variable is linearized by a basis function transformation. Each basis function can handle one sharp bend in the function. If more than one slope change is identified, several basis functions or an interaction may be required for their modeling.

The linearization renders the variable amenable to linear regression analysis, whereas the nonlinear form might be overlooked in a variable selection process.

The resulting linear combination of basis functions is then used in lieu of a nonlinear regression analysis. These functions are added to the linear combination of basis functions that constitutes the MARS model. For example, when PTSD in wave three is run against average cumulative reconstructed external dose of the respondent to ^{137}CS for women in Figure 1 or men in Figure 2, we can




Figure 1: Comparison of PTSD anxiety relationship before and after basis function transformation

observe that the relationships are nonlinear.

MARS generates sets of pairs of truncated spline transformations for each possible position on the domain of the regressor. They are generated in reflected pairs, always representing the positive part of the domain. These plots often appear as hockey sticks of various dimensions. But just as our patterns of male PTSD against cumulative external dose of $^{137}\text{Cesium}$ reveal piecewise patterns, these hockey stick transformations can serve to linearize delayed, latent, or threshold responses to exogenous variables. For these reasons, relationships that have such a shape may be linearized by MARS and made more amenable to linear OLS regression analysis than otherwise might be the case. Because this is the first empirical examination of this subject, we are partly in an exploratory mode. Therefore, we want to be sure that we explore as many aspects of the relationships that might exist that we can. In situations such as these MARS may provide an invaluable investigative tool to plumb the depths of relationships that could exist within our dataset.

After generating many of these regression splines from the radial basis functions, the pair that di-

minished the sum of squared errors the most was the pair that was incorporated into the model.[17, 287].

To guard against overfitting by the forward step-wise algorithm, backward pruning was also incorporated into the system, while the model undergoes a 10-fold cross-validation test. Alternatively, the user may opt for a testing segment to be reserved, and then a training segment to preclude that overfitting.

The nonparametric selection of basis function provided a different driver for the model building process that could be modulated at a later stage. It is that last stage where AutoMetrics is brought into the process.

9 Dose reconstruction of exposure to ^{137}CS

It is helpful to know that the ^{137}CS in the Ukraine was not in general very large. Natural background radiation is about 2.4 mSv/year. Depending upon where people were and what they ate, and the extent to which they were outdoors, the amount of radiation to which they were ex-

posed may well have been below the level of normal biological reactivity.

However, that was not always the case. Infant children under the age of 2-5, when their thyroids absorb more iodine from the air, were more affected than older youth whose physical growth had slowed down. When natural iodine uptake from the air had considerably subsided on the part of the youth of five or older, as it does in the natural life cycle, the danger subsided. Moreover, the farther away they were from the accident site, the less their health was threatened, unless the winds shifted direction and carried the radioactive plume to them. For some time, there was considerable uncertainty as to who and how much the health of some people were threatened. In the Ukraine the in area we sampled, the situation can be described by Figure 2.

Because our sample was a random one, many of our respondents lived beyond the reach of the exclusion zone, an area of approximately 30 km from Chernobyl. Consequently, the mean effective dose sustained by this sample may not reflect the condition of those who were seriously and substantially exposed.




Figure 2: Cumulative ^{137}CS dose in the Ukraine in microGrays

Table 1: **Deposition of ^{137}CS**

Measure of ^{137}CS	μGrays
minimum:	44
maximum:	26,600
mode:	800
median:	715
mean:	838
standard deviation:	1695
geometric mean:	550
geometric standard deviation:	2.4

We hope to test this hypothesis with the best possible regression model. What does that mean? We have to have the optimal covariates in the model to assure that the general unrestricted model (GUM) is congruent. We wish to minimize the possibility of specification error by increasing the chance of including relevant covariates in our model. We use MARS to point out potential relationships that we might have missed by automatically and systematically finding relationships in the data by recentering and transforming our variables.

10 Application of MARS and AutoMetrics to variable selection and model building

10.1 BSI positive symptoms models

10.1.1 Female positive symptoms models

10.1.2 Female positive symptoms model using AutoMetrics alone

Table 2 Modelling BSIPosymp by OLS-CS

The dataset is: gals.in7

The estimation sample is: 1 - 363

Dropped 3 observation(s) with missing values from the sample

	Coefficient	Std.Error	HACSE	t-HACSE	t-prob	Part.R ²
Constant	36.1449	7.460	6.939	5.21	0.0000	0.0741
age	0.567945	0.1656	0.1737	3.27	0.0012	0.0306
edu3	8.27144	2.726	3.183	2.60	0.0098	0.0195
edu4	-1.86104	5.139	4.521	-0.412	0.6809	0.0005
marrw13	3.87829	3.807	4.073	0.952	0.3416	0.0027
marrw16	-4.68635	11.63	7.866	-0.596	0.5517	0.0010
marrw24	32.7286	23.94	6.326	5.17	0.0000	0.0732
childw1	-2.08219	2.009	2.003	-1.04	0.2993	0.0032
emplw16	10.1987	3.583	3.277	3.11	0.0020	0.0278
emplw23	2.59385	4.633	4.253	0.610	0.5423	0.0011
emplw24	2.61391	24.74	7.494	0.349	0.7275	0.0004
depagw2	0.237815	0.07052	0.06819	3.49	0.0006	0.0346
anxagw1	0.146863	0.03891	0.04042	3.63	0.0003	0.0375
radfmw1	0.0553968	0.08792	0.07394	0.749	0.4542	0.0017
radhlw1	-0.180692	0.09119	0.08640	-2.09	0.0372	0.0127
radhlw3	0.234851	0.04998	0.05035	4.66	0.0000	0.0603
inc4w1	9.02537	5.808	6.853	1.32	0.1887	0.0051
inc1w3	10.2430	3.593	4.756	2.15	0.0320	0.0135
avgcumdosew1 U	4.72494	4.655	4.306	1.10	0.2733	0.0035
avgcumdosew2 U	4.50510	5.896	5.738	0.785	0.4329	0.0018
avgcumdosew3 U	-3.55804	4.255	3.947	-0.901	0.3680	0.0024
sigma	23.0765	RSS	180526.366			
R ²	0.365256	F(20,339) =	9.754	[0.000]**		
Adj.R ²	0.327808	log-likelihood	-1629.97			
no. of observations	360	no. of parameters	21			
mean(BSIPosymp)	86.6667	se(BSIPosymp)	28.1465			
When the log-likelihood constant is NOT included:						
AIC	6.33419	SC	6.56088			
HQ	6.42433	FPE	563.590			
When the log-likelihood constant is included:						
AIC	9.17207	SC	9.39876			
HQ	9.26221	FPE	9625.82			
Normality test:	Chi ² (2) =	58.204	[0.0000]**			
Hetero test:	F(28,329) =	1.1833	[0.2433]			
Hetero-X test:	F(73,284) =	0.87766	[0.7442]			
RESET23 test:	F(2,337) =	1.9412	[0.1451]			

10.1.3 Female model using AutoMetrics and MARS

Table 3 Modelling BSIPosymp by OLS-CS
The dataset is: gals.dta
The estimation sample is: 1 - 363

	Coefficient	Std.Error	HACSE	t-HACSE	t-prob	Part.R ²
BFps2f	1.77871	0.1518	0.2158	8.24	0.0000	0.1629
BFps3f	3.64609	0.07497	0.09200	39.6	0.0000	0.8182
BFps3	-3.44030	0.1022	0.1302	-26.4	0.0000	0.6667
BFps4	3.53564	0.07835	0.1000	35.3	0.0000	0.7816
BFps8w2	-0.638613	0.1687	0.1754	-3.64	0.0003	0.0366
BFps9	-1.70877	0.09559	0.1195	-14.3	0.0000	0.3692
BFps10	2.40125	0.1976	0.2177	11.0	0.0000	0.2585
BFps11	-1.96492	0.1354	0.1634	-12.0	0.0000	0.2930
BFps12	0.155755	0.04992	0.06037	2.58	0.0103	0.0187
BFps22aw1	-0.346626	0.2059	0.1572	-2.20	0.0281	0.0137
BFps28w2	-4.59650	2.512	1.921	-2.39	0.0173	0.0161
avgcumdosew1 U	0.188333	1.400	1.189	0.158	0.8742	0.0001
avgcumdosew2 U	1.86420	1.758	1.550	1.20	0.2300	0.0041
avgcumdosew3 U	-1.23718	1.263	1.287	-0.961	0.3371	0.0026

```

sigma          6.94119  RSS          16814.877
R^2            0.941297  F(13,349) =  430.5 [0.000]**
Adj.R^2        0.939111  log-likelihood  -1211.24
no. of observations      363  no. of parameters      14
mean(BSIPosymp)    86.5069  se(BSIPosymp)      28.1296
When the log-likelihood constant is NOT included:
AIC              3.91275  SC              4.06295
HQ              3.97245  FPE             50.0384
When the log-likelihood constant is included:
AIC              6.75063  SC              6.90083
HQ              6.81033  FPE             854.628B

```

```

// basis function legend
//-----
// 0 bf4m = max(0, 32 - BSIsoma)
// 1 BFps2f = max(0, phobanx - 2.03628e-00*)
// 2 BFps3f = max(0, WHPer + 3.14933e-007)
// 3 BFps3 = max(0, WHPer - 33.7)
// 4 BFps4 = max(0, 33.7 - WHPer)
// 5 BFps8w2 = max(0, 5 - depagw2)
// 6 BFps9 = max(0, 3.852327e-007)
// 7 BFps10= max(0, BSIPsysc - 4)
// 8 BFps11 = max(0, 17- BSIPar)
// 9 BFps12 = max(0, MiPTSD-44)
// 10 BFps22aw1= max(0, 3-pillw1)
// 11 BFps28w2 = max(0, sepaw2 + 5.41039e-010)

```

```

Normality test:  Chi^2(2) = 18.105 [0.0001]**
Hetero test:     F(25,337) = 1.8159 [0.0108]*
Hetero-X test:   F(97,265) = 2.0224 [0.0000]**
RESET23 test:    F(2,347) = 1.5619 [0.2112]

```

10.2 Male Models for BSI Positive symptoms

Particular problems plague these men during wave two with significant effects. They include air and water pollution in both waves one and two (airw1 and airw2), and self-reported wave two depression and Chernobyl related health threat to oneself (radhlw2). However, the only dose-positive symptom impact is evident at wave one. The model as measured by adjusted $R^2=0.458$ is a respectable. But even more important is that only one of the model assumptions is in violation.

10.2.1 Male model with AutoMetrics Alone

Table 4: EQ(13) Modelling BSIPosymp by OLS-CS
The dataset is: guys.dta
The estimation sample is: 1 - 340

	Coefficient	Std.Error	HACSE	t-HACSE	t-prob	Part.R^2
Constant	76.0152	4.446	4.030	18.9	0.0000	0.5328
educ4	10.5398	4.830	5.188	2.03	0.0431	0.0131
educ7	-13.1150	9.002	4.358	-3.01	0.0028	0.0282
marrw12	-15.9041	7.535	3.753	-4.24	0.0000	0.0544
marrw21	-12.1409	2.996	3.263	-3.72	0.0002	0.0425
marrw24	-19.2302	10.39	4.196	-4.58	0.0000	0.0631
marrw33	-6.72192	2.293	2.648	-2.54	0.0116	0.0202
childw2	-3.42953	1.581	1.863	-1.84	0.0666	0.0107
emplw25	-14.6481	4.954	4.564	-3.21	0.0015	0.0320
occ6w1	-14.7849	8.032	5.540	-2.67	0.0080	0.0223
occ8w1	-3.04011	2.796	3.050	-0.997	0.3196	0.0032
occ7w3	9.52459	2.796	3.381	2.82	0.0052	0.0248
inc1w1	4.51683	3.031	2.836	1.59	0.1122	0.0081
inc3w2	-2.83038	2.105	2.050	-1.38	0.1684	0.0061
inc4w2	-10.3680	6.052	5.953	-1.74	0.0826	0.0096
inc1w3	-1.23529	3.226	3.836	-0.322	0.7476	0.0003
inc4w3	11.6291	5.642	4.657	2.50	0.0130	0.0196
airw1	-0.0538877	0.03236	0.03894	-1.38	0.1674	0.0061
airw2	-0.108320	0.04605	0.04958	-2.18	0.0296	0.0151
airw3	0.0562605	0.04006	0.04158	1.35	0.1770	0.0058
radchw3	-0.0375368	0.03117	0.02909	-1.29	0.1979	0.0053
radhlw2	0.224384	0.03387	0.04215	5.32	0.0000	0.0833
illw3	6.08420	1.103	1.337	4.55	0.0000	0.0623
depagw1	0.0908117	0.04081	0.06095	1.49	0.1372	0.0071
depagw2	0.269824	0.07069	0.1049	2.57	0.0106	0.0208
avgcumdosew1 U	1.18106	1.284	0.5961	1.98	0.0484	0.0124
avgcumdosew2 U	-1.43189	4.240	2.289	-0.626	0.5320	0.0013
avgcumdosew3 U	0.560894	3.831	2.229	0.252	0.8015	0.0002
sigma	17.0257	RSS	90441.298			
R^2	0.501136	F(27,312) =	11.61	[0.000]**		
Adj.R^2	0.457965	log-likelihood	-1431.64			
no. of observations	340	no. of parameters	28			
mean(BSIPosymp)	74.9618	se(BSIPosymp)	23.1256			
When the log-likelihood constant is NOT included:						
AIC	5.74822	SC	6.06354			
HQ	5.87386	FPE	313.748			
When the log-likelihood constant is included:						
AIC	8.58609	SC	8.90142			
HQ	8.71174	FPE	5358.65			
Normality test:	Chi^2(2) =	35.773	[0.0000]**			
Hetero test:	F(39,300) =	1.3885	[0.0694]			
RESET23 test:	F(2,310) =	0.55927	[0.5722]			

10.2.2 Male model for Positive symptoms using AutoMetrics + MARS

Table 5 Modelling BSIPosymp by OLS-CS

The dataset is guys.in7

The estimation sample is: 1 - 340

	Coefficient	Std.Error	HACSE	t-HACSE	t-prob	Part.R ²
depagw1	0.0422038	0.01124	0.01496	2.82	0.0051	0.0245
BFps1f	1.01014	0.3436	0.5369	1.88	0.0608	0.0110
BFps2f	3.77415	0.4887	1.001	3.77	0.0002	0.0429
BFps4	0.800758	0.09437	0.1582	5.06	0.0000	0.0748
BFps14b	3.54876	0.5376	1.033	3.44	0.0007	0.0359
BFbsidep11	-0.0370855	0.01675	0.01997	-1.86	0.0643	0.0108
BFbsidep12	1.66140	0.1198	0.1356	12.3	0.0000	0.3214
BFbsidep13	2.09106	0.1552	0.2106	9.93	0.0000	0.2373
BFbsidep14	2.30043	0.1575	0.1948	11.8	0.0000	0.3055
BFptsd2a	0.0397467	0.01259	0.01901	2.09	0.0373	0.0136
BF5sociso3	-2.24763	0.3732	0.5600	-4.01	0.0001	0.0484
BFdep1	0.776016	0.08013	0.1486	5.22	0.0000	0.0792
emplw11	22.1433	4.671	10.71	2.07	0.0395	0.0133
emplw12	16.3055	3.151	8.316	1.96	0.0508	0.0120
emplw13	16.7090	3.211	8.284	2.02	0.0445	0.0127
emplw16	16.9370	3.194	8.240	2.06	0.0407	0.0132
emplw22	1.71709	0.6972	0.6371	2.70	0.0074	0.0224
occ5w2	2.15340	1.128	1.186	1.82	0.0704	0.0103
occ4w3	1.41793	0.8934	0.6351	2.23	0.0263	0.0155
I:180	-35.9714	5.501	3.896	-9.23	0.0000	0.2120
avgcumdosew1 U	-0.678928	0.3604	0.3830	-1.77	0.0772	0.0098
avgcumdosew2 U	0.552981	1.172	1.380	0.401	0.6889	0.0005
avgcumdosew3 U	-0.0585932	1.055	1.196	-0.0490	0.9609	0.0000

```

sigma          4.85845  RSS          7482.62731
log-likelihood -1007.98
no. of observations      340  no. of parameters      23
mean(BSIPosymp)    74.9618  se(BSIPosymp)    23.1256
When the log-likelihood constant is NOT included:
AIC          3.22669  SC          3.48570
HQ          3.32989  FPE          25.2013
When the log-likelihood constant is included:
AIC          6.06456  SC          6.32358
HQ          6.16777  FPE          430.424

```

```

//-----
// Basis function legend for positive symptom analysis
//0  bf4m = max(0, 32 - BSIsoma)
//1  BFps1f = max(0, bf4m - 1.19546e-007)
//2  BFps2f = max(0, phobanx - 2.03628e-008)
//3  BFps4 = max(0, 33.7 - WHPer)
//4  BFbsidep11 = max(0, 65.06 - WHPsleep)
//5  BFbsidep12 = max(0, BSIsoc - 5)
//6  BFbsidep13 = max(0, BSIsanx - 5)
//7  BFbsidep14 = max(0, BSIsips - 4)
//8  BFps14b = max(0, 11-BSIphanx)
//10 BFptsd2a = max(0, fdferw2 - 7.17707e-007)
//11 BF5sociso3 = max(0, 25-BSIsoma)
//12 BFdep1 = max(0, WHPer + 3.14933e-007)

```

```

Normality test:  Chi^2(2) = 18.319 [0.0001]**
Hetero test:    F(37,301) = 6.5782 [0.0000]**
RESET23 test:   F(2,315) = 20.030 [0.0000]**

```

When we add basis functions to the model, we observe the results in Table 5. We find no evidence to support a dose-mental health relationship. We observe a tendency of the radial basis functions to dominate the output, with 47.8% of the variables selected are now basis functions. Because we now need variable labels to be able to facilitate an interpretation of the output, we provide a basis function index at the base of the model output.

The assumption violation rate increases again from 33.3% to 100%.

Many other factors appear to be related to general mental health on the part of the males, including depression in 1986 (depagw1 and BF5socio3), somaticism (BFps1f), lack of sleep (BFbsidep11), anxiety (BFbsidep13 and BFbsi14b), interpersonal sensitivity (BFbsidep14), emotional reaction (BFps4 and BFdepx1), with some obsessive-compulsiveness (BSIdep12) included.

10.3 PTSD models

To test the hypothesis that cumulative dose predicts PTSD among men and women, we examine the addition of truncated regression splines does to the model of dose-PTSD relationship.

First we want to graphically examine the relationship in Figure 3. Not only do we see that the relationship changes over time, but that at first it is characterized by an abrupt turn downward in wave one. In later waves, it more or less straightens out but retains a slight upward slope.

First we run our baseline model before adding the truncated spline transformations. Unlike our previous findings, we find that dose effect relationship seem to be associated with PTSD on the part of the male respondents. This association remains through all three waves.

We find comfort in believing that AutoMetrics has kept those covariates which were significantly related to the endogenous variable and which therefore provide us with the good control against specification error, at least insofar as the Ramsay reset test would have us believe. However, two of the regression assumptions are not fulfilled.

10.4 Male PTSD models

Socioeconomic status appears to play a significant role in PTSD for males. Being employed full-time during wave one (emplw12) is positively associated with PTSD. But part-time employment dur-




Figure 3: Male PTSD against Cumulative Effective Dose of ^{137}CS over three waves

ing wave three (emplw33) is significantly negatively associated with PTSD. For some reason, homemaking and caregiving becomes positively associated with male PTSD in wave three.

Many of these men have experienced some kind of catastrophe in wave three (cataw3) and they report illnesses in waves one and three (illw1 and illw3). There is a significant yet inverse relationship between their belief in the proportion of pollution due to Chornobyl (radchw3) in wave three. They are aware of the hazardous effects of radiation (efradw2) and remain fearful of eating radioactively contaminated food in wave three (fdferw3).

These are men who exhibit significant emotional reactions (WHPer), somaticism (BSIsoma), paranoia (BSIpar), and in wave two they report anxiety (anxagw2). For some reason the direction of significant dose-PTSD relationship switches from inverse to direct and then fades into relative statistical non-significance in wave three ($p=0.074$).

10.4.1 Male PTSD model using AutoMetrics alone

Table 6 EQ(20) Modelling MiPTSD by OLS-CS

The dataset is:/guys.dta

The estimation sample is: 1 - 340

Dropped 1 observation(s) with missing values from the sample

	Coefficient	Std.Error	HACSE	t-HACSE	t-prob	Part.R ²
emplw12	1.74095	0.8400	0.7437	2.34	0.0199	0.0170
emplw25	9.57283	3.406	6.409	1.49	0.1363	0.0070
emplw33	-7.60494	1.677	1.891	-4.02	0.0001	0.0487
occ7w2	-8.96825	3.580	6.720	-1.33	0.1830	0.0056
occ7w3	5.29819	1.689	1.802	2.94	0.0035	0.0266
cataw3	15.0846	3.670	5.033	3.00	0.0029	0.0276
illw1	2.82366	0.9845	1.149	2.46	0.0145	0.0188
illw3	1.56530	0.4444	0.4377	3.58	0.0004	0.0389
shfamw1	-0.00984147	0.009452	0.009251	-1.06	0.2882	0.0036
suchrw3	0.0144091	0.009844	0.009279	1.55	0.1214	0.0076
fdferw3	0.133797	0.01621	0.01669	8.02	0.0000	0.1690
efradw2	0.0607720	0.01242	0.01325	4.59	0.0000	0.0624
radchw1	0.0134653	0.01282	0.01344	1.00	0.3170	0.0032
radchw3	-0.0395825	0.01386	0.01374	-2.88	0.0042	0.0256
WHPer	0.112901	0.03244	0.03768	3.00	0.0029	0.0276
LBSItotal	7.31530	0.4480	0.4154	17.6	0.0000	0.4954
BSIsoma	0.502610	0.09646	0.1127	4.46	0.0000	0.0592
BSIips	-0.337637	0.2135	0.1954	-1.73	0.0850	0.0094
BSIpar	0.393373	0.1629	0.1462	2.69	0.0075	0.0224
anxagw2	0.0638708	0.02325	0.02468	2.59	0.0101	0.0207
avgcumdosew1 U	-1.03625	0.4535	0.3685	-2.81	0.0052	0.0244
avgcumdosew2 U	3.53115	1.577	1.736	2.03	0.0427	0.0129
avgcumdosew3 U	-2.80259	1.423	1.566	-1.79	0.0744	0.0100
sigma	6.1492	RSS		11948.787		
log-likelihood	-1084.84					
no. of observations	339	no. of parameters		23		
mean(MiPTSD)	47.174	se(MiPTSD)		11.9189		
When the log-likelihood constant is NOT included:						
AIC	3.69808	SC		3.95766		
HQ	3.80152	FPE		40.3781		
When the log-likelihood constant is included:						
AIC	6.53596	SC		6.79554		
HQ	6.63940	FPE		689.636		
Normality test:	Chi ² (2)	=	7.3566	[0.0253]*		
Hetero test:	F(40,298)	=	1.6234	[0.0133]*		
RESET23 test:	F(2,314)	=	1.0049	[0.3673]		

10.4.2 Male PTSD model using AutoMetrics + MARS

By adding basis functions, the model becomes considerably less parsimonious. In fact, the number of parameters increases to 65, although there is a huge decline in the residual sums of squares from 11948.787 to 6783.70162. The Schwartz criterion is used as an arbiter, this actually rises from 3.958 to about 4.16 with the addition of the new variables. The only basis function that was added in this case was a linear combination of the dependent variable, which we subsequently deleted from the variable candidate pool.

10.5 Male PTSD model with AutoMetrics + MARS- cont'd

Table 7 EQ(25) Modelling MiPTSD by OLS-CS

The dataset is: guys.dta

The estimation sample is: 1 - 340

Dropped 9 observation(s) with missing values from the sample

	Coefficient	Std.Error	HACSE	t-HACSE	t-prob	Part.R ²
emplw12	7.69107	2.819	2.420	3.18	0.0017	0.0366
occ7w2	-5.04297	1.831	1.736	-2.91	0.0040	0.0308
occ7w3	10.3099	2.226	2.254	4.57	0.0000	0.0729
cataw3	8.28971	4.078	2.440	3.40	0.0008	0.0416
illw1	5.17788	0.9376	0.9342	5.54	0.0000	0.1035
fdferw3	0.0302069	0.03236	0.03779	0.799	0.4248	0.0024
efradw2	0.0238300	0.01675	0.01587	1.50	0.1343	0.0084
radchw1	0.0169080	0.01422	0.01504	1.12	0.2619	0.0047
radchw3	-0.0567978	0.01438	0.01427	-3.98	0.0001	0.0562
lBSItotal	4.20572	1.033	0.9615	4.37	0.0000	0.0671
BSIips	-0.911879	0.2381	0.2334	-3.91	0.0001	0.0543
anxagw2	0.0497071	0.02337	0.02327	2.14	0.0336	0.0169
emplw13	5.89476	2.982	2.564	2.30	0.0223	0.0195
emplw16	6.08702	2.940	2.568	2.37	0.0185	0.0207
occ1w1	-2.59177	1.095	0.9997	-2.59	0.0101	0.0246
occ2w2	-3.66632	1.256	1.134	-3.23	0.0014	0.0378
occ3w2	-3.72039	1.623	1.487	-2.50	0.0129	0.0230
occ4w2	-3.96241	1.373	1.072	-3.70	0.0003	0.0489
occ8w2	-2.27052	1.193	1.002	-2.27	0.0243	0.0189
occ1w3	10.7594	2.171	2.255	4.77	0.0000	0.0788
occ2w3	11.5444	2.323	2.244	5.14	0.0000	0.0905
occ3w3	10.5016	2.460	2.383	4.41	0.0000	0.0680
occ4w3	9.37193	2.393	2.131	4.40	0.0000	0.0678
occ5w3	11.9117	2.347	2.450	4.86	0.0000	0.0816
occ6w3	6.08384	3.680	5.175	1.18	0.2408	0.0052
inc2w3	-2.02479	0.9227	0.9335	-2.17	0.0310	0.0174
inc3w3	-2.38019	0.9749	1.008	-2.36	0.0189	0.0205
cataw1	-1.43900	0.9247	0.8904	-1.62	0.1072	0.0097
cataw2	7.34793	3.360	2.425	3.03	0.0027	0.0334
dvcew3	-2.54128	1.302	1.335	-1.90	0.0581	0.0134
illw2	1.16389	0.5908	0.5776	2.02	0.0449	0.0150
movew2	4.36285	0.8182	0.6689	6.52	0.0000	0.1379
shjobw1	0.0436225	0.01294	0.01375	3.17	0.0017	0.0365
shjobw2	-0.0396988	0.01321	0.01286	-3.09	0.0022	0.0346

Table 7 -- continued...

	Coefficient	Std.Error	HACSE	t-HACSE	t-prob	Part.R ²
shhlw2	0.0223050	0.01419	0.01356	1.65	0.1011	0.0101
shfincw2	-0.0219229	0.01411	0.01400	-1.57	0.1187	0.0091
shfincw3	0.0136628	0.01356	0.01175	1.16	0.2459	0.0051
shhousw1	-0.0436495	0.01381	0.01440	-3.03	0.0027	0.0334
shhousw3	0.0403242	0.01213	0.01228	3.28	0.0012	0.0389
shrelaw2	0.0375747	0.01576	0.01483	2.53	0.0119	0.0236
shrelaw3	-0.0429053	0.01358	0.01398	-3.07	0.0024	0.0342
suprtw3	0.00665292	0.008794	0.008390	0.793	0.4285	0.0024
sufamw1	0.0500712	0.01839	0.02032	2.46	0.0144	0.0223
suchrw2	-0.0200943	0.008790	0.009018	-2.23	0.0267	0.0183
fdferw2	0.0793413	0.03196	0.04034	1.97	0.0502	0.0143
kmacc	0.0823621	0.01387	0.02051	4.02	0.0001	0.0572
injothr	1.59261	0.7767	0.8111	1.96	0.0506	0.0143
kmwork	-0.0861868	0.01378	0.02045	-4.21	0.0000	0.0626
polprw3	0.0358676	0.01554	0.01554	2.31	0.0217	0.0196
airw2	-0.0413224	0.01281	0.01355	-3.05	0.0025	0.0338
radw1	0.0322737	0.01101	0.01106	2.92	0.0038	0.0310
radhlw1	-0.0385425	0.01545	0.01560	-2.47	0.0141	0.0224
radhlw3	0.0455350	0.01831	0.01870	2.44	0.0155	0.0218
healthef	0.0176544	0.01307	0.01555	1.14	0.2572	0.0048
icdxcnt	-0.395851	0.2306	0.2331	-1.70	0.0906	0.0107
HP2work	1.58433	0.8974	0.7985	1.98	0.0483	0.0146
HP2hmcare	-2.48808	0.9763	0.8842	-2.81	0.0053	0.0289
HP2pbfhm	1.92806	1.379	1.448	1.33	0.1842	0.0066
BSIposymp	0.382404	0.05583	0.05262	7.27	0.0000	0.1656
BSIdep	-0.617556	0.1875	0.1736	-3.56	0.0004	0.0454
BSIphanx	-1.03931	0.2448	0.2323	-4.47	0.0000	0.0700
BSIhos	-0.425587	0.1974	0.1798	-2.37	0.0186	0.0206
avgcumdosew1 U	0.730692	0.7812	0.5074	1.44	0.1510	0.0077
avgcumdosew2 U	-3.98866	2.462	1.707	-2.34	0.0202	0.0201
avgcumdosew3 U	2.47899	1.851	1.423	1.74	0.0827	0.0113
sigma	5.05001	RSS	6783.70162			
log-likelihood	-969.505					
no. of observations	331	no. of parameters	65			
mean(MiPTSD)	47.0242	se(MiPTSD)	11.7831			
When the log-likelihood constant is NOT included:						
AIC	3.41291	SC	4.15955			
HQ	3.71070	FPE	30.5107			
When the log-likelihood constant is included:						
AIC	6.25079	SC	6.99743			
HQ	6.54858	FPE	521.107			
Normality test:	Chi ² (2) =	0.97273	[0.6149]			
Hetero test:	F(104,226)=	1.1353	[0.2170]			
RESET23 test:	F(2,264) =	12.918	[0.0000]**			

With this model, the hypothesis that dose predicts PTSD is not supported by the data for males at wave one and three. The only apparent statistically significant association appears during wave

two (avgcumdosew2, $b = -3.989$, $se = 2.462$, $t\text{-prob} = 0.0202$). That wave, covering the years of 1987 through 1996, was one of substantial economic privation and difficulty, during which other stresses and strains may have become more salient in the minds of these men.

What were the significant covariates? One group consisted of socio-demographic characteristics— such as being employed full, part, or unemployed in wave one (emplw12 emplw13 and emplw16, respectively)— were somehow positive associated with Civilian PTSD in wave three.

Some aspects of socio-economic status appear to be related to PTSD in wave three. Having been a professional or administrator in wave one was negatively related to PTSD. In wave 2, employment in administrative support or technical sales (occ2w2), the protective services (occ3w2), precision production, mechanical, or crafts fields (occ3w2), or being a student (occ8w2) at the time were negatively associated with PTSD in wave three, as measured by the civilian PTSD scale score. By wave 3, such employment — along with employment as a factory laborer (occ5w3) or in agriculture, forestry, trapping, or logging (occ6w3) with the exception of being

a student, were positively associated with PTSD. Income sufficiency with resources enough to meet basic necessities plus a little left over (inc3w3) is negatively associated with PTSD. It appears as if socio-economic adequacy may be important.

Environmental factors may have played a role as well. Males suffering from PTSD were concerned about the distance of their residence and workplace from the scene of the accident (kmacc and kmwork). By wave two, their concerns about air and water pollution (airw2) were inversely associated with PTSD. The farther the distance from the accident site the less the PTSD. They were concerned about the Chornobyl related threats to their own health (radhlw3) in wave three, whereas at first any concern they had was inversely associated with PTSD (radhlw1). They were people whose reported illnesses count in waves one (illw1) and two (illw2) were significantly associated with PTSD. Their fear of eating radioactively contaminated food was of borderline statistical significance in wave two (fdferw2 $p=0.0502$), which became statistically nonsignificant later.

PTSD appears to be positively related to having observed a catastrophic event in wave two or three

(cataw2 or cataw3). A belief in the proportion of radioactively contaminated area is directly related to PTSD. Stresses and hassles from job related matters were in wave one directly associated with PTSD but switched to a significant inverse related relationship in wave two (shjobw1 and shjobw2). Stresses and hassles due to housing matters were inversely related in wave one but directly related in wave three (shhousw1 and shhousw3). Stresses and hassles relating to relationships went from significantly positive to significantly negative in waves two and three (shrelaw2 and shrelaw3).

Forms of support were occasionally related to PTSD in wave3. Family support was significant only in wave one (sufamw1). Chornobyl survivor support was negatively related to PTSD in wave two (suchrw2).

These were individuals who exhibited general mental health dysfunctionality (BSIposymp) whose health problems impacted their work (HP2work). Both of these traits were significantly positively related to PTSD in wave three. There was a significant positive association with PTSD between the log of total BSI. There was a borderline significance in the relationship of having injured others

and PTSD ($p=0.0506$). Their reported anxiety in wave two was statistically significant with PTSD in wave three ($\text{anxagw2}, b = .049, p = 0.034$).

They also exhibited a number of significant inverse relationships between psychological symptomatology and PTSD. These included depression (BSIdep), phobic anxiety (BSIphanx), hostility (BSIhos), interpersonal sensitivity (BSIips), and the impact of health issues on home care cooking, and repairs (HP2hmcare). It is possible that the economic difficulties exacerbated the PTSD problems in wave two so that there appeared to be a dose-PTSD male relationship at that time.

10.6 Female PTSD models using AutoMetrics alone

Before turning to the female PTSD models, it behooves us to graph the relationships observed between PTSD and average cumulative dose over the three waves, shown in Figure 4.

Wave one appears to be less linear in functional form than the others, so we may have more need of basis functions there, but the other two relationships appear to be fairly linear and without the need of transformation.

We can examine the female baseline model ex-




Figure 4: Female PTSD over three waves

plaining PTSD, for our hypothesis test of stressors, and then of buffers relating to the PTSD. In the baseline model, we observe no statistically significant relationship between cumulative external dose as measured by avgcumdose in waves one through three, and PTSD for women, measured by the Civilian PTSD scale score.

For the women, the baseline model has 55 parameters. Two-thirds of the first page of output consist in socio-economic attributes. How does socio-economic status relate to PTSD? Some of the largest coefficients indicate salient aspects of

socio-economic status related to PTSD. Being single (marrw21) in wave two 2 can be positively related to it. Income insufficiency (inc1w3) or almost having such insufficiency (inc2w3) in wave 3 are significantly related to the PTSD. Having a child in wave one is significantly inversely related to it. With respect to occupational status, working outdoors (occ6w1) is significantly related to PTSD in wave two (farming, logging, trapping, etc).

Other stressors are salient. Accidents (accdw1) in wave1 are significantly positively related to female PTSD in wave three. Stresses and hassles from relationships (shrelaw1) during wave one are significantly positively related to PTSD. Catastrophes in wave three are positive related to PTSD for women. General mental dysfunctionality as measured by the BSI positive symptom scale is significantly positively related to PTSD. Self-reported depression (depagw1 and depagw2) and anxiety (anxagw1 and anxagw2) may be statistically related as well. They seem to be significantly related but the signs of their parameter estimate change in opposite directions as we move from wave one to wave two. Health problem interference with interests and hobbies (HP2inthob) is significantly

positively associated with PTSD on the part of the female respondents. Consumption of pain pills (pillw2) in wave two seems to be statistically related to PTSD. The geodesic distance of the residence from the accident site (havmil) is also found to be statistically significantly related to PTSD. Also, the stresses and hassles from the job and from relationships appears to be significantly positively related to female PTSD.

Buffers to PTSD are also evident in this model. Having a child in wave one (childw1) for a woman is inversely related to PTSD. Being an MD (occ8w3) is also inversely related to PTSD. Working outdoors in agriculture, logging, forestry, are other jobs inversely related to PTSD (occ6w3). Health issues impacting one's sex life seem to be inversely related to PTSD for women (HP2sxlife). The stresses and hassles from one's health are inversely related to PTSD for women.

Supports also have a role to play. Family support(sufamw2) in wave two is significantly related as is Chornobyl survivor support in wave one (suchrw1). The belief that a large proportion of pollution is due to Chornobyl (radchw3) is significantly positively related to female PTSD.

Table 8 EQ(29) Modelling MiPTSD by OLS-CS

The dataset is: gals.dta

The estimation sample is: 1 - 363

Dropped 7 observation(s) with missing values from the sample

	Coefficient	Std.Error	HACSE	t-HACSE	t-prob	Part.R ²
marrw12	8.24237	4.931	4.806	1.72	0.0873	0.0096
marrw21	12.3299	1.805	1.792	6.88	0.0000	0.1355
marrw22	8.47335	3.703	2.766	3.06	0.0024	0.0301
marrw23	6.23052	1.614	1.725	3.61	0.0004	0.0414
childw1	-1.26565	0.8791	0.7999	-1.58	0.1146	0.0082
childw2	2.55125	0.9794	0.8653	2.95	0.0034	0.0280
emplw13	2.94446	2.047	1.774	1.66	0.0980	0.0090
emplw32	3.52670	1.876	2.169	1.63	0.1050	0.0087
emplw33	4.18612	7.971	1.777	2.36	0.0191	0.0180
occ2w1	2.77946	1.582	1.613	1.72	0.0860	0.0097
occ4w1	-0.606786	2.170	1.844	-0.329	0.7424	0.0004
occ6w1	13.1978	4.550	2.815	4.69	0.0000	0.0678
occ6w2	-9.92894	4.473	2.773	-3.58	0.0004	0.0407
occ8w2	-5.30301	2.045	2.187	-2.42	0.0159	0.0191
occ8w3	-9.21187	8.387	2.808	-3.28	0.0012	0.0344
inc1w3	8.22857	1.680	1.812	4.54	0.0000	0.0639
inc2w3	8.85438	1.391	1.337	6.62	0.0000	0.1267
inc3w3	7.71199	1.413	1.372	5.62	0.0000	0.0947
deaw1	1.16721	0.5685	0.6472	1.80	0.0723	0.0107
deaw2	-0.631489	0.5636	0.5662	-1.12	0.2656	0.0041
dvcew3	-5.08077	1.884	1.790	-2.84	0.0048	0.0260
sepaw2	-5.42709	3.268	2.621	-2.07	0.0392	0.0140
sepaw3	7.34660	2.257	2.257	3.26	0.0013	0.0339
accdw1	8.14050	2.308	1.849	4.40	0.0000	0.0603
accdw3	4.52347	1.349	1.181	3.83	0.0002	0.0463
cataw1	-2.97149	1.501	1.410	-2.11	0.0359	0.0145
cataw3	9.48043	5.937	5.320	1.78	0.0757	0.0104
shjobw1	0.0459037	0.01897	0.02220	2.07	0.0395	0.0140
shfamw2	0.0173869	0.02005	0.02067	0.841	0.4010	0.0023
shhlw1	-0.0632661	0.01947	0.02554	-2.48	0.0138	0.0199
shhlw3	0.0113486	0.01642	0.01671	0.679	0.4976	0.0015
shfincw2	-0.0151884	0.02226	0.02365	-0.642	0.5212	0.0014
shfincw3	-0.0204790	0.01681	0.01948	-1.05	0.2941	0.0036
shrelaw1	0.0579751	0.01556	0.01523	3.81	0.0002	0.0458
sufamw3	0.0354931	0.01329	0.01453	2.44	0.0151	0.0194
suchrw1	0.0885848	0.04288	0.04238	2.09	0.0374	0.0143
pillw2	0.246420	0.1558	0.1249	1.97	0.0493	0.0127
pillw3	-0.132645	0.09234	0.09077	-1.46	0.1450	0.0070
injselfr	1.48063	1.100	1.147	1.29	0.1978	0.0055

Table 8 - continued...

explanatory var	Coefficient	Std.Error	HACSE	t-HACSE	t-prob	Part.R ²
radchw1	-0.00608758	0.01568	0.01719	-0.354	0.7234	0.0004
radchw3	0.0455101	0.01787	0.01961	2.32	0.0210	0.0175
WHPer	0.0124274	0.03620	0.04239	0.293	0.7696	0.0003
HP2pbfhm	-2.00175	1.695	1.999	-1.00	0.3176	0.0033
HP2sxlife	-5.74679	1.371	1.523	-3.77	0.0002	0.0450
HP2inthob	7.84715	1.583	2.314	3.39	0.0008	0.0367
BSIposymp	0.292820	0.02081	0.02386	12.3	0.0000	0.3328
havmil	0.00610682	0.001950	0.002717	2.25	0.0253	0.0165
depagw1	-0.0666916	0.02328	0.02283	-2.92	0.0037	0.0275
depagw2	0.109470	0.03546	0.03460	3.16	0.0017	0.0321
anxagw1	0.0455276	0.01835	0.01819	2.50	0.0128	0.0203
anxagw2	-0.0720609	0.02882	0.03571	-2.02	0.0445	0.0133
avgcumdosew1 U	3.03671	1.669	1.928	1.57	0.1164	0.0081
avgcumdosew2 U	2.25711	2.073	2.156	1.05	0.2960	0.0036
avgcumdosew3 U	-2.93719	1.507	1.815	-1.62	0.1067	0.0086
sigma	7.70878	RSS		17946.4376		
log-likelihood	-1202.94					
no. of observations	356	no. of parameters		54		
mean(MiPTSD)	49.6208	se(MiPTSD)		12.0886		
When the log-likelihood constant is NOT included:						
AIC	4.22359	SC		4.81136		
HQ	4.45739	FPE		68.4392		
When the log-likelihood constant is included:						
AIC	7.06146	SC		7.64923		
HQ	7.29527	FPE		1168.91		
Normality test:	Chi ² (2) =	4.9117	[0.0858]			
Hetero test:	F(81,272) =	1.1388	[0.2219]			
RESET23 test:	F(2,300) =	37.383	[0.0000]**			

10.6.1 Female PTSD model using AutoMetrics and MARS

When we mix AutoMetrics with MARS to test the female PTSD hypothesis, we get an elaborate rather than a parsimonious model. The hypothesis that dose is directly associated with PTSD measured by the Civilian Mississippi scale is consistent with parameter estimates for cumulative dose of ¹³⁷CS in mGys (avgcumdosew1 and avgcumdosew3, both of which are statistically significant at the 0.05 level).

Table 9 EQ(32) Modelling MiPTSD by OLS-CS

The dataset is:gals.dta

The estimation sample is: 1 - 363

Dropped 5 observation(s) with missing values from the sample

	Coefficient	Std.Error	HACSE	t-HACSE	t-prob	Part.R ²
marrw10	2.40743	3.127	2.103	1.14	0.2533	0.0044
marrw12	8.70104	3.872	4.456	1.95	0.0518	0.0126
marrw13	3.79369	1.166	1.149	3.30	0.0011	0.0351
marrw21	3.82439	1.410	1.323	2.89	0.0041	0.0272
marrw25	7.03501	3.217	2.954	2.38	0.0179	0.0186
marrw32	-4.37927	2.753	1.800	-2.43	0.0155	0.0194
marrw35	-5.37582	1.928	2.308	-2.33	0.0205	0.0178
childw1	-0.595462	0.5982	0.7546	-0.789	0.4307	0.0021
emplw13	3.04047	1.861	1.736	1.75	0.0810	0.0102
emplw15	5.87467	3.790	2.599	2.26	0.0245	0.0168
emplw22	2.45999	1.227	1.047	2.35	0.0194	0.0181
emplw25	-3.04349	2.065	2.118	-1.44	0.1518	0.0069
emplw33	8.96577	7.031	2.166	4.14	0.0000	0.0542
occ3w1	-5.70773	1.423	1.598	-3.57	0.0004	0.0409
occ6w1	5.11728	2.547	2.332	2.19	0.0290	0.0159
occ2w2	2.73149	1.220	1.465	1.86	0.0632	0.0115
occ7w2	5.13912	1.805	1.523	3.38	0.0008	0.0367
occ8w2	-3.99304	1.859	1.748	-2.28	0.0231	0.0172
occ8w3	-7.08976	6.967	1.908	-3.72	0.0002	0.0442
inc1w3	2.59772	1.277	1.311	1.98	0.0485	0.0130
inc2w3	2.52167	0.9265	0.9327	2.70	0.0073	0.0239
deaw2	-0.776765	0.5023	0.5270	-1.47	0.1416	0.0072
dvcew1	3.21334	5.055	2.772	1.16	0.2473	0.0045
dvcew2	-6.65500	2.146	1.942	-3.43	0.0007	0.0378

Continued on the next page...

Table 9 continued from previous page ...

	Coefficient	Std.Error	HACSE	t-HACSE	t-prob	Part.R ²
sepaw1	-11.6999	5.853	5.004	-2.34	0.0200	0.0180
sepaw3	5.04579	1.756	1.970	2.56	0.0109	0.0215
accdw1	8.29547	1.976	2.055	4.04	0.0001	0.0517
accdw3	4.28951	1.165	1.373	3.12	0.0020	0.0316
cataw3	12.6758	5.373	5.723	2.21	0.0275	0.0161
illw3	1.44130	0.4337	0.4956	2.91	0.0039	0.0275
movew3	1.88537	2.146	1.747	1.08	0.2813	0.0039
shjobw1	0.0507772	0.01648	0.01829	2.78	0.0058	0.0251
shjobw3	0.00812849	0.01277	0.01383	0.588	0.5572	0.0012
shhlw1	-0.0444529	0.01734	0.02059	-2.16	0.0317	0.0153
shfincw1	0.0448605	0.01388	0.01257	3.57	0.0004	0.0408
shfincw3	-0.0395446	0.01394	0.01599	-2.47	0.0139	0.0201
suprtw2	-0.0222859	0.009940	0.009618	-2.32	0.0212	0.0176
sufamw3	0.00922673	0.01160	0.009733	0.948	0.3439	0.0030
pillw3	-0.114727	0.06201	0.04760	-2.41	0.0165	0.0191
airw1	0.0250637	0.01327	0.01270	1.97	0.0494	0.0129
radchw1	-0.0600910	0.01499	0.01546	-3.89	0.0001	0.0481
radchw3	0.0758247	0.01561	0.01648	4.60	0.0000	0.0661
WHPsleep	0.0585702	0.01677	0.01846	3.17	0.0017	0.0326
WHPsociso	-0.0184077	0.02557	0.02705	-0.680	0.4967	0.0015
HP2sxlife	-4.72120	1.209	1.364	-3.46	0.0006	0.0385
HP2vacatn	5.48797	1.353	1.482	3.70	0.0003	0.0439
BSIposymp	0.208521	0.01950	0.02045	10.2	0.0000	0.2579
depagw1	-0.0678127	0.02029	0.02263	-3.00	0.0030	0.0292
depagw2	0.100628	0.03216	0.03028	3.32	0.0010	0.0356
anxagw1	0.0612168	0.01615	0.01665	3.68	0.0003	0.0432
anxagw2	-0.157812	0.03253	0.03638	-4.34	0.0000	0.0592
BFptsd5	0.266666	0.04240	0.03959	6.74	0.0000	0.1318
BFptsd2a	0.277405	0.03833	0.03685	7.53	0.0000	0.1593
BFptsd4a	0.0842508	0.04848	0.04852	1.74	0.0835	0.0100
BFptsd5a	-0.0270958	0.01449	0.01617	-1.68	0.0949	0.0093
BFptsdw33	0.0974791	0.03504	0.03758	2.59	0.0100	0.0220
avgcumdosew1 U	4.28430	1.459	1.855	2.31	0.0216	0.0175
avgcumdosew2 U	2.65230	1.847	1.800	1.47	0.1417	0.0072
avgcumdosew3 U	-3.54964	1.341	1.492	-2.38	0.0180	0.0186
sigma	6.68318	RSS	13354.8046			
log-likelihood	-1155.8					
no. of observations	358	no. of parameters	59			
mean(MiPTSD)	49.6117	se(MiPTSD)	12.0553			
When the log-likelihood constant is NOT included:						
AIC	3.94871	SC	4.58824			
HQ	4.20305	FPE	52.0259			
When the log-likelihood constant is included:						
AIC	6.78658	SC	7.42611			
HQ	7.04093	FPE	888.574			
Normality test: Chi ² (2) = 5.0543 [0.0799]						
Hetero test: F(88,267) = 1.4085 [0.0201]*						
RESET23 test: F(2,297) = 1.4425 [0.2380]						

Basis function legend:

BFptsd5 = max(0, 70-fdferw2)
BFptsd4a = max(0, 40 - depagw3)
BFptsd5a = max (0, airw2-10)
BFpstdw33 = max(0, anxagw3)

To summarize the female analysis, we will note some salient significant stressors and buffers for PTSD among women in several domains—that of socio-economic status (SES), that of major negative life events, that of daily stresses and hassles, environmental factors, and psychological sequelae. Using the partial R^2 as a form of beta weight, we see that within the domain of SES, being married in wave 1 (marrw13), being divorced in wave 2 (marrw25), unemployed in wave 3 (emplw35), and having a Ph.D. in wave 2 (occ7w2) are major stressors relating to PTSD. Other SES stressors are income insufficiency for basic needs or borderline income sufficiency for basic needs in wave three (inc1w3 and inc2w3).

Prominent socio-economic status buffers are being retired in waves 2 and 3 (emplw25 and emplw35) and being a medical doctor in waves two and three (occ8w2 and occ8w3).

Among the major negative life events, several salient stressors appear. Some of the worst are getting separated in wave one (sepaw1) and getting a divorce in wave two, that time of economic tribulation (dvcew2). However, accidents in wave one (accdw1) and three (accdw3) also emerge as

substantial stressors.

Within the domain of daily stresses and hassles, financial ones in wave 1 are significantly positively related to PTSD. The reception of Chornobyl support in wave 2 and the consumption of pain pills in wave three (pillw3) not surprisingly appear to work as buffers.

As for environmental or contextual effects, in wave 1 the percent belief air and water pollution (airw1) is dangerous is related to PTSD, as is the proportion of pollution due to Chornobyl is high in wave 3 (radchw3) appears to be significantly negatively related to PTSD, although there was a belief that the air and water pollution was dangerous in wave 1 (airw1) and wave 2 (BFptsd5a), whereas by wave three the direction of this relationship reversed itself, so that it became a significantly positive in wave 3.

Fears of eating contaminated food appear to be significantly positively related to PTSD in wave 2 (BFptsd5).

Psychological symptoms appeared in depression and anxiety in waves one and two (depagw1 and depage2), where they went from inverse to direct over those two waves. However, anxiety went from

positive to negative as people became used to the situation from wave one to wave two (anxagw1 and anxagw2). The basis functions reveal that depression and anxiety in wave three were related to PTSD as well.

These effects impacted the sleep, sex life, and vacation plans of the female respondents such that even general mental dysfunction became significantly positively related to PTSD. When the nature of the situation is considered, these women may not seem so unreasonable, after all.

11 Conclusion

11.1 Exploration of new dimensions

Does MARS help us explore dimensions we might otherwise have ignored? Sometimes it does and sometimes it does not. For the men, the proportion of variance explained tends to increase, but for the females it does not. In our study of the BSI positive symptom response, males exhibit an increase in the number of factors extracted, but females do not. Females, to the contrary, exhibit a slight decline in the number of factors extracted. In our PTSD models, the same pattern appears.

The result may vary with the nature of the data.

11.2 Are MARS models well-behaved?

MARS models tend to violate more assumptions than AutoMetrics built models and MARS basis function generation supervised by AutoMetrics does not guarantee well-behaved residuals. Sometimes MARS generated basis functions if selected by AutoMetrics proliferate assumption violations and sometimes they do not. The results at this juncture are not conclusive, although there is tentative evidence that MARS is motivated by optimizing fit and not fulfilling regression assumptions. Therefore the user must be careful about his use of MARS.

MARS interaction generation fails to include all component parts of the interaction and actually may prune some of them out if it improves the overall fit. The resulting model may be misspecified according to conventional statistical theory.

11.3 Can MARS capture nonlinear relationships ?

MARS can by adding basis functions approximate a nonlinear relationship between the dependent

variable and the other variables in the model. However, it may try to do this by building one of its incomplete and unbalanced interaction vectors. This may not be what the user wants so he must be careful about such developments.

11.4 Content validity Tables

Does MARS improve content validity? The answer is sometimes and not always. If the dimensionality of the explanatory variables increases, then the content validity increases. But the use of MARS may actually decrease the dimensionality of the model, as shown among women with respect to their Positive symptom scale scores and their Chornobyl PTSD scores in Tables 2 and 3 below. For male respondents the application of MARS appeared to increase dimensionality but did not necessarily improve fit.

Did MARS always improve the fit? If we examine in table 2 immediately above the proportion of common variance explained we find that among male BSI positive symptoms, it did. However, among females, the proportion of common variance explained declined. As for the PTSD, the application of MARS as a front end generator of

Table 2: Construct validity assessment of BSI positive symptom models

	AutoMetrics alone	MARS + AutoMetrics
Male model		
nfactors	4	6
Number of singletons	0	0
% common variance explained:	87	94
number of regression predictors	28	23
Number of regression assumption violations	1	2
Female model		
nfactors	3	2
Number of singletons	0	0
% common variance explained	92	87
Number of regression assumption violations	0	0
number of regression predictors	21	14
Number of regression assumption violations	1	2

basis functions did improve proportion of variance explained among men but not in women. If the dimensionality of the predictor variable set actually decline slightly among women. Further research needs to be done to be sure, but if these few examples can serve as a guide, the value added by MARS is mostly to capture nonlinearities in relevant relationships.

Table 3: Construct validity assessment of PTSD models

	AutoMetrics	MARS+AutoMetrics
Male model		
nfactors	4	15
Number of singletons	0	0
% common variance explained:	85	88
number of regression predictors	23	65
number of regression assumption violations	1	3
Female model		
nfactors	10	9
Number of singletons	0	0
% common variance explained	86	84
number of regression predictors	54	59
number of regression assumption violations	1	1

11.5 Advantages of MARS

- automated
- fast
- builds additive model for you

11.6 Disadvantages of MARS

- adding them together to get a function
- no prima facie clear translation available sometimes
- you will need a translation table
- only emphasizes goodness of fit
- does not test regression assumptions
- interaction terms are not conventional

11.7 Caveats

- MARS may provide an optimal fit at the expense of violating the assumptions of regression analysis
- does not test regression assumptions
- interaction terms are not conventional in that they do not include the component terms conventionally required for proper interaction specification.

References

- [1] Havenaar, J.M, de Wilde, E.J., Van Den Bout, J., Drottz-Sjoberg, B.M. and Van Den Brink, W. (2003) Perception of risk and subjective health among victims of the Chernobyl disaster *Social Science and Medicine*, 569-572.
- [2] Slovic, P. (2000) *The Perception of Risk* London:Earthscan Publications Ltd.
- [3] Bromet, E. J. and Litcher, L. (2002) *Psychological response of mothers of young children to the Three Mile Island and Chernobyl*

Nuclear Plant accidents one decade later in
In J.M. Havenaar, J. Cwikel and E.J. Bromet
(Eds.) *Toxic Turmoil: Psychological and So-
cietal Consequences of Ecological Disasters*
NY:Kluwer Academic/Plenum Publishers.

- [4] Dew, M.A., Bromet, E.J., Schulburg, H.C.,
Dunn, L.O. and Parkinson, D.K. (1987) Mental
health effects of the Three Mile Island nuclear
reactor restart *American Journal of Psychia-
try*, 144, 1074-1077.
- [5] Page, W.G., Babyleva, A.E., Naboka, M.V.
and Shestopalov, M.V. (1995) *Environmental
health policies in Ukraine after the Chernobyl
accident* Policy Studies Journal, 23, 141-151.
- [6] Michal, F., Grigor, K.M. Negro-Vilar, A. and
Skakkebaek, N.E. (1993) *Impact of environ-
ment on reproductive health* *Environmental
Health Perspectives*, 101 (Suppl. 2) 159-167.
- [7] Amirkhan, J.H. and Auyeung, B. (2007) Cop-
ing with stress across the life-span: Absolute
versus relative changes in strategy *Journal of
Applied Developmental Psychology*, 28, 298-
317.
- [8] Purvis-Roberts, K.L., Werner, C.A., and

- Frank, I. (2007) Perceived risks from radiation and nuclear testing near Semipalatinsk, Kazakhstan: A comparison between physicians, scientists and the public. *Risk Analysis*, 291-302.
- [9] Hendry, Sir D.F. and Richard, J-F. (1983) The econometric analysis of economic time series (with discussion). *International Statistical Review*, 51, 111-163.
- [10] Hendry, Sir D.F., and Krolzig, H-M. (2001) Automatic Econometric Model Selection using Pc-GETS London, UK: Timberlake Consultants, Ltd..
- [11] Hunt, S.M., McEwen, J. and McKenna, S.P. (1985) Measuring health status: A new tool for clinicians and epidemiologists *Journal of the Royal College of General Practitioners*, 35, 185-188.
- [12] Perez Foster, R., Yaffee, R.A., Borak, T., and Tierney, K. (2010) Modeling nuclear disaster risk Poster presented at the National Science Foundation, Human and Social Dynamics Grantees Conference, Sept. 26-27, 2010, Arlington.

- [13] Derogatis, L.R. (1993) Brief Symptom Inventory: Procedures Manual. MN:National Computers Systems.
- [14] Perez Foster, R. and Goldstein, M.F. (2007) Chernobyl disaster sequelae in recent immigrants to the US from the former Soviet union *Journal of Immigrant Health*, 9, 115-124.
- [15] Castle, J.L., Doornik, J.A., and Hendry, D.F. 2011 Evaluating Automatic Model Selection *Journal of Time Series Econometrics*, Vol.3, 1, 1-31.
- [16] Friedman, Jerome H. 1990 Multivariate Adaptive Regression Splines *Stanford Linear Accelerator Center Publication 4960-Rev*.
- [17] Hastie, T., Tibsharani, R. and Friedman, J. 2001 *The Elements of Statistical Learning* New York, N.Y.: Springer, 115-163.
- [18] Harrell, Jr., Frank 2001 *Regression Modeling Strategies* New York, N.Y.: Springer, 18-24.
- [19] Lewis, P.A.W. and Stevens, J. G. 1991 Non-linear Modeling of Time Series Using Multivariate Adaptive Regression Splines *Journal of*

the American Statistical Association, Vol.86,
No.416, 864-877.

- [20] Alina V. Brenner, Mykola D. Tronko, Maureen Hatch, Tetyana I. Bogdanova, Valery A. Oliynik, Jay H. Lubin, Lydia B. Zablotska, Valery P. Tereschenko, Robert J. McConnell, Galina A. Zamotaeva, Patrick OKane, Andre C. Bouville, Ludmila V. Chaykovskaya, Ellen Greenebaum, Ihor P. Paster, Victor M. Shpak, Elaine Ron (2007) I-131 Dose Response for Incident Thyroid Cancers in Ukraine Related to the Chornobyl Accident <http://ehp03.niehs.nih.gov/article/info:doi/10.1289/ehp.1002674>.
- [21] Marsh, L.C.and Cormier, David R. 2001 Spline Regression Models Sage Publications
- [22] Chernobyl Forum (2003-2005) (2–5) Chernobyl Legacy: Health, Environmental and Socio-Economic Impacts: Recommendations to the Governments of Belarus, Russian Federation and Ukraine, Second Revised Edition.
- [23] MARS User Guide (2001) San Diego, Ca: Salford Systems, Ince., 2, 15.
- [24] Selvin, Steve (2011) *Statistical tools for Epi-*

demiological Research New York: Oxford University Press, 257-309.